

The Air4EU project is financed by the 6th framework programme of the European commission.

Bruce Denby

Norwegian Institute for Air Research (NILU), P.O. Box 100, 2027 Kjeller, Norway



Summary

This case study examines the use of basic data assimilation methods and how they can improve the spatial assessment of air quality on the European scale. In previous studies Horálek et al. (2005) produced interpolated maps of PM₁₀ based on annual statistics for Europe by combining the unified EMEP model with observations. As part of this case study the assimilation methodology has been further developed to make use of Bayesian statistics to combine observational and model fields and to calculate the annual mean and number of exceedance days based on the daily mean statistics, rather than annual values. This poster looks at the differences that occur when applying the assimilation method to annual or daily mean statistics. The maps represent the rural background concentrations of PM₁₀ averaged over a 25 x 25 km grid.



Assimilation methodology

The assimilation method used here combines a linearly fitted model regression field with an interpolated observational field using kriging. The two fields are combined based on minimisation of the variance using Bayesian theory. Uncertainty of the regression model is estimated from the normalised root mean square error (NRMSE). The uncertainty field is then made by multiplying the model field with the NRMSE. The uncertainty of the observational field is determined from the kriged variance field used in the kriging interpolation. Uncertainty is also assessed using the cross validated RMSE. Observational data has been retrieved from the AirBase and EMEP databases.

Annual mean for PM₁₀ 2003





Original model field as calculated using the unified EMEP model for the year 2003. Also shown are the observations with availability > 75%, circles. Crosses indicate stations also used in the daily statistics but with total availability < 75%





The assimilated map is created by the weighted combination of linear model regression with kriged interpolation fields. The weighting, according to Bayesian theory, minimises the uncertainty in the final map. Annual mean values have been used



Number of exceedance (NOE) days for PM₁₀ 2003





Original model exceedance field as calculated using the unified EMEP model for the year 2003. Also shown are the observations for availability > 75%, circles. Crosses indicate stations also used in the daily statistics but with total availability < 75%





The assimilated map is created by the weighted combination of linear model regression with kriged interpolation fields. The weighting, according to Bayesian theory, minimises the uncertainty in the final map. Annual mean values have been used.





The assimilated map is created using assimilation on a daily basis, rather than annual.



The map is created by assimilating the daily mean concentrations. The number of exceedances are calculated based on these daily mapped exceedances, rather than by mapping the annual number of exceedances.



The difference between the annual and daily mean assimilation fields. The annually based method gives slightly higher values in polluted regions. The difference is smaller than the uncertainty.



The difference between the annual and daily assimilated exceedance fields. The difference is of the same order as the uncertainty but are in some areas significant.

Model

Assimilation: annual statistics

Assimilation: daily statistics

Difference: annual - daily

Conclusion

- Basic data assimilation methods have been used to create maps of the annual mean PM₁₀ and of the number of daily exceedances over the limit value for PM₁₀ on the European scale. These methods significantly improve the spatial mapping for both these indicators.
- Comparison of the data assimilation method applied to annual or daily statistics indicates that:
 - There are only small differences between the annual and daily assimilation on mean concentrations of PM₁₀ and these are less than their estimated uncertainty.
 - There are significant differences between the annual and daily assimilation of PM₁₀ when calculating the number of exceedance days.
- It is recommended to use annual statistics when calculating annual means.
- It is recommended to use daily statistics when calculating daily exceedances.
- Both the cross validation RMSE and general uncertainty maps are useful in ascertaining the uncertainty of the assimilated field. The methodology for producing uncertainty maps requires further research.
- The inclusion of uncertainty maps is a very useful indicator of the uncertainty in the assessment maps and is highly recommended for any air quality mapping.